

Recycling and Reuse Technology Transfer Center

Recycling and Reuse Technology Transfer Center



<http://www.rrttc.uni.edu>

**Gas phase reactions of conjugated dienes as a model for pyrolysis
Submitted to Combustible Flame 1995**

Publication: 1995 – 042

Curtiss Hanson, Department of Chemistry, and Michaela Rich, RRTTC, University of Northern Iowa

Student authors: M. Hammer, Chemistry Department, University of Northern Iowa

95-042

Gas phase Reactions of Conjugated Diene Systems as a Model for Pyrolysis

Michaela L. Rich, Michelle Hammer, Tim Burrell, Curtiss D. Hanson, Department of Chemistry, University of Northern Iowa, Cedar Falls, IA 50614

The pyrolysis of styrene-butadiene rubber (SBR) produces large molecular weight aromatic and aliphatic hydrocarbons through secondary reactions in the gas phase. Fourier transform ion cyclotron resonance mass spectrometry (FT-ICR) was used to analyze the products produced by the gas phase reactions that occur during the pyrolysis of SBR tires. Conjugated diene systems were used on the FT-ICR as models for these reactions in order to better understand the process through which secondary pyrolysis products are formed. Deuterium labeling experiments were utilized to determine the mechanism for the model reactions. Relative rates of reaction and product distribution of the conjugated diene systems will be shown to be consistent with those from the SBR tire pyrolysis.

9-042

Gas phase Reactions of Conjugated Diene Systems as a Model for Pyrolysis

Michaela L. Rich, Michelle Hammer, Tim Burrell, Curtiss D. Hanson, Department of Chemistry, University of Northern Iowa, Cedar Falls, IA 50614

The pyrolysis of styrene-butadiene rubber (SBR) produces large molecular weight aromatic and aliphatic hydrocarbons through secondary reactions in the gas phase. Fourier transform ion cyclotron resonance mass spectrometry (FT-ICR) was used to analyze the products produced by the gas phase reactions that occur during the pyrolysis of SBR tires. Conjugated diene systems were used on the FT-ICR as models for these reactions in order to better understand the process through which secondary pyrolysis products are formed. Deuterium labeling experiments were utilized to determine the mechanism for the model reactions. Relative rates of reaction and product distribution of the conjugated diene systems will be shown to be consistent with those from the SBR tire pyrolysis.